

U.S. FISH AND WILDLIFE SERVICE SPECIES ASSESSMENT AND LISTING PRIORITY ASSIGNMENT FORM

Scientific Name:

Etheostoma spilotum ssp.

Common Name:

Kentucky arrow Darter

Lead region:

Region 4 (Southeast Region)

Information current as of:

04/18/2013

Status/Action

Funding provided for a proposed rule. Assessment not updated.

Species Assessment - determined species did not meet the definition of the endangered or threatened under the Act and, therefore, was not elevated to the Candidate status.

New Candidate

Continuing Candidate

Listing Priority Number (LPN) Change

Former LPN: 3

New LPN: 2

Candidate Removal

Taxon is more abundant or widespread than previously believed or not subject to the degree of threats sufficient to warrant issuance of a proposed listing or continuance of candidate status

Taxon not subject to the degree of threats sufficient to warrant issuance of a proposed listing or continuance of candidate status due, in part or totally, to conservation efforts that remove or reduce the threats to the species

Range is no longer a U.S. territory

Insufficient information exists on biological vulnerability and threats to support listing

Taxon mistakenly included in past notice of review

Taxon does not meet the definition of "species"

Taxon believed to be extinct

Conservation efforts have removed or reduced threats

More abundant than believed, diminished threats, or threats eliminated.

Petition Information

Non-Petitioned

Petitioned

90-Day Positive:

12 Month Positive:

Did the Petition request a reclassification?

For Petitioned Candidate species:

Is the listing warranted(if yes, see summary threats below)

To Date, has publication of the proposal to list been precluded by other higher priority listing?

Explanation of why precluded:

We find that the immediate issuance of a proposed rule and timely promulgation of a final rule for this species has been, for the preceding 12 months, and continues to be, precluded by higher priority listing actions (including candidate species with lower LPNs). During the past 12 months, the majority our entire national listing budget has been consumed by work on various listing actions to comply with court orders and court-approved settlement agreements; meeting statutory deadlines for petition findings or listing determinations; emergency listing evaluations and determinations; and essential litigation-related administrative and program management tasks. We will continue to monitor the status of this species as new information becomes available. This review will determine if a change in status is warranted, including the need to make prompt use of emergency listing procedures. For information on listing actions taken over the past 12 months, see the discussion of Progress on Revising the Lists, in the current CNOR which can be viewed on our Internet website (<http://endangered.fws.gov/>).

Historical States/Territories/Countries of Occurrence:

- **States/US Territories:** Kentucky
- **US Counties:** Breathitt, KY, Clay, KY, Harlan, KY, Knott, KY, Lee, KY, Owsley, KY, Perry, KY, Wolfe, KY
- **Countries:** United States

Current States/Counties/Territories/Countries of Occurrence:

- **States/US Territories:** Kentucky
- **US Counties:** Breathitt, KY, Clay, KY, Harlan, KY, Knott, KY, Lee, KY, Leslie, KY, Owsley, KY, Wolfe, KY
- **Countries:**Country information not available

Land Ownership:

Extant populations of the Kentucky arrow darter occur in watersheds that are privately owned and publicly owned (U.S. Forest Service, Daniel Boone National Forest [DBNF] and Robinson Forest, University of Kentucky [UK]). The DBNFs ownership is typically fragmented but is most concentrated within the South Fork Kentucky River basin in Clay, Leslie, and Owsley Counties. Robinson Forest is located in Breathitt and Knott Counties.

Lead Region Contact:

ARD-ECOL SVCS, Erin Rivenbark, 706 613-9493, erin_rivenbark@fws.gov

Lead Field Office Contact:

KY ESFO, Michael Floyd, 502-695-0468, mike_floyd@fws.gov

Biological Information

Species Description:

The Kentucky arrow darter, *Etheostoma spilotum* Gilbert, is a rather large darter reaching 116 millimeters (mm) (4.6 inches [in]) total length (TL) (Kuehne and Barbour 1983, p. 71; Etnier and Starnes 1993, p. 523). It has a slender body, elongated snout, large mouth, and virtually scaleless head. The species ground color is straw yellow to pale greenish, and the back is crossed by 5 to 7 weak dorsal saddles, some of which may fuse with the 8 to 11 vertical lateral blotches (Kuehne and Barbour 1983, p. 71; Etnier and Starnes 1993, p. 523). Anterior blotches are generally oval with pale centers. Posteriorly, blotches extend ventrally almost to the midline and may resemble the letters N, W, U, or V. A dark vertical bar occurs at the base of the caudal fin, sometimes separated by two distinct spots. The belly is pale (Kuehne and Barbour 1983, p. 71).

During the spawning season, breeding males exhibit vibrant coloration. Most of the body is blue-green in color, with scattered scarlet spots and scarlet to orange vertical bars laterally; the vertical bars can be connected ventrally by an orange belly stripe (Etnier and Starnes 1993, p. 523). The spinous dorsal fin exhibits a blue-green central band and a scarlet marginal band. The soft dorsal and caudal fins are speckled with scarlet blotches or bands, and the anal and pelvic fins are blue-green to black. Females remain pale straw yellow with grayish markings (Etnier and Starnes 1993, p. 523).

Taxonomy:

The Kentucky arrow darter was described from the Kentucky River basin (Sturgeon Creek, Owsley County) as *Etheostoma nianguae spilotum* (Gilbert 1887, pp. 53-54). Bailey (1948, pp. 80-84) redescribed the species, placing it and its closest relatives, *P. nianguae* (Gilbert and Meek) and *P. sagitta* (Jordan and Swain), in a new subgenus, *Litocara*. Subsequent to this, Bailey et al. (1954, pp. 109-164) and Bailey and Gosline (1955, pp. 1-44) synonymized (combined taxonomically) *Poecilichthys* with the genus *Etheostoma* and *Litocara* with the subgenus *Oligocephalus*, in which *E. spilotum* and its relatives were regarded as a species group (group of closely related species within a genus that are grouped because they are morphologically similar and share a common ancestry). Kuehne and Bailey (1961, pp. 1-5) evaluated new material for all three members of *Oligocephalus* and determined that the group consisted of two species, *E. nianguae* (Gilbert and Meek) and *E. sagitta* (Jordan and Swain), and three forms. *Etheostoma nianguae* was distinctive morphologically and confined to the lower Osage River system in Missouri. *Etheostoma sagitta* was recognized as a polytypic species (represented by more than one subspecies), consisting of *E. s. sagitta*, an endemic form to the upper Cumberland River system, and *E. sagitta spilotum* (Kentucky arrow darter), an

endemic form to the upper Kentucky River system. The subgenus *Litocara* was later resurrected (reinstated taxonomically) by Page and Whitt (1973, pp. 611-623) to include *E. nianguae* Gilbert and Meek and *E. sagitta* (Jordan and Swain). Subsequently, the Kentucky arrow darters subspecific status was supported by Kuehne and Barbour (1983, p. 71), Page (1983, p. 59), and Etnier and Starnes (1993, p. 523).

Thomas (2008, p. 6) questioned the polytypic status of *E. sagitta* by arguing that (1) the two subspecies were distinguishable based on scale size and the development of the lateral line (see note below), (2) the two subspecies existed in allopatry (separate ranges with no overlap), (3) the two subspecies lacked intergrades (intermediate forms), and (4) unpublished genetic data (mitochondrial DNA) suggested evolutionary independence of Kentucky and Cumberland basin populations (with no recent genetic exchange). Based on these analyses and additional morphological and genetic evidence presented by Thomas and Johansen (2008, p. 46), the two arrow darter subspecies have been elevated to species rank (Page and Burr 2011, p. 569; Eschmeyer 2013). The Cumberland arrow darter, *E. sagitta* (Jordan and Swain) is restricted to the upper Cumberland River basin, and the Kentucky arrow darter, *E. pilotum* Gilbert, is restricted to the upper Kentucky River basin.

*Note: Cumberland and Kentucky arrow darters are indistinguishable based on general appearance, including pigment pattern and breeding color; however, the two species are separable based on various scale counts. Thomas (2008, p. 6) examined specimens of both species and determined that the Kentucky arrow darter had lateral scale counts of 62 or fewer in 88% of individuals examined (vs. 63 or more in 94% of Cumberland arrow darters), pored lateral scale counts of 50 or fewer in 79% of individuals examined (vs. 51 or more in 91% of Cumberland arrow darters), and caudal peduncle scale counts of 22 or fewer in 72% of individuals examined (vs. 23 or more in 83% of Cumberland arrow darters). These differences reflect a trend toward larger scale size and a more weakly developed lateral line (a faint line of sense organs extending from the gill cover to the tail) in the Kentucky arrow darter.

Habitat/Life History:

During the most recent range-wide surveys (Thomas 2008, p. 6; USFWS 2010, pp. 1-13), Kentucky arrow darters were found in pools or transitional areas between riffles and pools (runs and glides) in moderate-to high-gradient streams. Individuals were usually associated with bedrock, boulder, and cobble substrates and occasionally observed around woody debris. Stream widths ranged from 1.5 to 20 meters (m) (5 to 66 feet [ft]), and depths at which individuals were captured ranged from 10 to 45 centimeters (cm) (4 to 18 in). During 2007 and 2008, Thomas (2008, p. 6) observed Kentucky arrow darters in streams ranging in size from first to third order, with 60 percent occurring in second order streams. The majority (72 percent) of these streams were in watersheds draining an area of 20 square kilometers (km²) (7.7 square miles [mi²]) or less. Many of these habitats, especially those in first order reaches, can be intermittent in nature. Lotrich (1973, p. 394) observed riffle habitats in Clemons Fork (Breathitt County) that were completely dry by late summer. These habitats continued to support arrow darters, but fishes were crowded into isolated pools once drying occurred.

Male darters establish territories over riffles from March to May, where they are quite conspicuous in water 5 to 15 cm (2 to 6 in) deep (Kuehne and Barbour 1983, p. 71). Males fan out a depression in the substrate and defend these sites vigorously. Initial courtship behavior involves rapid dashes, fin-flaring, nudging, and quivering motions by the male followed by similar quivering responses of the female, who then precedes the male to the nest. The female partially buries herself in the substrate, is mounted by the male, and spawning occurs (Etnier and Starnes 1993, p. 523). It is assumed that the male continues to defend the nest until the eggs have hatched. Bailey (1948) described collected females as bulging with eggs in April, probably the peak spawning period. Lowe (1979) studied the biology of *E. sagitta* and determined that the peak spawning period was during April when water temperatures reached 13°C (55°F).

Young arrow darters can reach 50 mm TL by the end of the first year (Lotrich 1973, p. 384-385; Lowe 1979), and one-year olds are generally sexually mature and participate in spawning with older age classes (Etnier

and Starnes 1993, p. 523). Lotrich (1973, p. 384) indicated mean length at age 2 of about 65 mm (2.6 in) and was unable to differentiate between older age classes (age 3+). Lowe (1979) reported four age classes, but growth was variable after age 1.

Lotrich (1973, p. 381) reported that Kentucky arrow darters captured in 1967 and 1968 from Clemons Fork fed primarily on mayflies, specifically the families Heptageniidae (genus *Stenonema*) and Baetidae. Mayflies comprised 77 percent of identifiable food items (420 of 542 items) in 57 arrow darter stomachs. Large arrow darters (individuals over 70 mm [2.8 in] TL) appeared to specialize on small crayfish, as 7 of 8 stomachs contained crayfish ranging in size from 11 to 24 mm (0.4 to 0.9 in). Lotrich (1973, p. 381) considered this to be noteworthy since stomachs of small arrow darters (less than 70 mm [2.8 in]) and stomachs of other darter species did not contain crayfish. He suggested that larger arrow darters were utilizing a different energy source, thus removing themselves from direct competition for food with other fishes in first and second order streams. This would allow these larger individuals to exploit an abundant food source and survive in extreme headwater habitats. Other arrow darter food items reported by Lotrich (1973, p. 381) and Etnier and Starnes (1993, p. 523) included larval blackflies (family Simuliidae) and midges (Chironomidae), with lesser amounts of caddisfly larvae, stonefly nymphs, and beetle larvae. Etnier and Starnes (1993, p. 523) reported that juvenile arrow darters feed on microcrustaceans and dipteran larvae.

Common associates of the Kentucky arrow darter include creek chub (*Semotilus atromaculatus*), central stoneroller (*Camptostoma anomalum*), white sucker (*Catostomus commersonii*), emerald darter (*Etheostoma baileyi*), rainbow darter (*E. caeruleum*), fantail darter (*E. flabellare*), and Johnny darter (*E. nigrum*) (Kuehne 1962, p. 609; Lotrich 1973, p. 380; Thomas 2008, p. 7). Within first-order or headwater reaches of these stream systems, the species is most commonly associated with creek chub, fantail darter, rainbow darter, and Johnny darter.

Historical Range/Distribution:

The Kentucky arrow darters historical distribution was limited to the upper Kentucky River system in eastern Kentucky (Kuehne and Bailey 1961, pp. 3-4; Kuehne 1962, pp. 608-609; Lotrich 1973, p. 380; Branson and Batch 1983, pp. 1-15; Burr and Warren 1986, p. 316; Ray and Ceas 2003, pp. 1-15). Its distribution spanned portions of five subbasins: Red River (Rockbridge Fork of Swift Camp Creek), Sturgeon Creek, South Fork Kentucky River, Middle Fork Kentucky River, and North Fork Kentucky River (Thomas 2008, p. 3).

Current Range Distribution:

The Kentucky arrow darter continues to occupy portions of the upper Kentucky River basin in eastern Kentucky, including the five sub-basins listed above; however, recent surveys by Thomas (2009, pp. 3-6) and the U.S. Fish and Wildlife Service (USFWS 2009, pp. 1-4; 2010, pp. 1-13) revealed that the Kentucky arrow darter has disappeared from portions of its range. The species was observed at only 34 of 68 historic streams (50 percent) and 45 of 100 historic sites (46 percent) during surveys completed from 2007 to 2010 (Thomas 2009, pp. 3-6; USFWS 2009, pp. 1-4; USFWS 2010, pp. 1-13). In 2010, additional surveys were initiated by the KFO within the Kentucky arrow darters historic range but in streams lacking previous records for the subspecies (USFWS 2010, pp. 1-13). A total of 14 new streams were surveyed across the basin, but no Kentucky arrow darters were observed.

A synopsis of the Kentucky arrow darters current range is provided below and is arranged by sub-basin. The current number of streams and sites known to support Kentucky arrow darters in each sub-basin is listed parenthetically after the sub-basin name. The species currently occupies 35 streams within the upper Kentucky River basin.

Red River (1/1). Greenberg and Steigerwald (1981, p. 37) discovered a previously unknown population of Kentucky arrow darters in Rockbridge Fork, a tributary of Swift Camp Creek, Red River basin in Wolfe County. Rockbridge Fork is located within the Daniel Boone National Forest (DBNF), Cumberland District

(Red River Gorge area). Surveys by KDFWR in 2007 produced 10 individuals (representing at least 2 age classes) from an approximate 200-m (0.12-mi) sampling reach on Rockbridge Fork (Thomas 2008, p. 4). Based on survey results and observations of habitat conditions at the site, Thomas (2008, p. 4) concluded that the population was stable. The KFO resurveyed this reach in March 2011, collecting five individuals (representing two age classes). Portions of the sampling reach had been degraded by sedimentation (sand and silt), but overall habitat conditions were suitable for the subspecies.

Sturgeon Creek (2/2). Gilbert (1887, pp. 53-54) described the Kentucky arrow darter based on specimens collected from Little Sturgeon Creek near Travellers Rest, Owsley County. Based on those results and other collections since that time, the Kentucky arrow darter was known from eight historic sites and five streams within the Sturgeon Creek basin. Thomas (2008, pp. 3-4) surveyed seven of eight historic sites and three new sites (Duck Fork, Rowlette Branch, and Travis Creek) within the basin. Arrow darters were observed at only two sites in two streams - Wild Dog Creek (1 individual) and Travis Creek (2 individuals). Habitats within Cooperas Cave Branch appeared suitable for arrow darters, but none were observed. Thomas (2008, p. 4) reported that much of the basin had been cleared for agricultural and residential development, with denuded riparian zones, siltation, and eutrophication identified as potential stressors. Travis Creek represented a new stream and collection site for the subspecies.

South Fork Kentucky River (17/19). Some of the best remaining populations of Kentucky arrow darters are found within this basin. The majority of occupied streams within the basin are located on the DBNF (Redbird District) in Clay, Leslie, and Owsley Counties. These streams include the Buffalo Creek watershed in Owsley County, and several small, first- and second-order tributaries of the Red Bird River in Clay and Leslie Counties (Thomas 2008, p. 4). These streams are characterized by relatively intact riparian zones with little or no human development, high gradients with abundant riffles, cool temperatures, low conductivity (near baseline conditions, less than 100 μ S), and stable channels with clean cobble/boulder substrates. Thomas (2008, p. 4) observed some oil and gas development and logging activity in these watersheds, but the activities did not appear to be adversely affecting these streams. Thomas (2008, p. 4) and USFWS (2009, pp. 1-4) surveyed 31 of 36 historic sites in the South Fork Kentucky River basin and observed Kentucky arrow darters at 19 of those sites. Recent surveys by the U.S. Forest Service (USFS) demonstrated that robust populations of Kentucky arrow darters continue to occupy portions of the Laurel Fork basin (Buffalo Creek watershed) in southeastern Owsley County (Brandt, pers. comm., 2011). Surveys on Laurel Fork and Cortland Branch yielded an average of 32 Kentucky arrow darters per 600-m (1,968-ft) sampling reach.

In contrast to streams within the DBNF (Redbird District), the remaining watersheds in the South Fork Kentucky River basin (e.g., Pawpaw Creek, Lower Buffalo Creek, Buck Creek, Cow Creek, Island Creek, Sexton Creek, and Goose Creek) have more residential and agricultural development, especially within their stream valleys (Thomas 2008, p. 4). Riparian zones tend to be narrower with less canopy cover, channel substrates are composed of smaller particles, siltation is more prevalent, and stream conductivity is higher (>160 μ S). Four streams within these basins, Horse Creek (Clay County), Laurel Creek (Clay County), Lower Buffalo Creek (Lee and Owsley Counties), and Sexton Creek (Clay and Owsley Counties), have been placed on Kentucky's 303d list of impaired waters (KDOW 2008, pp. 65-101) because they were not supportive of the Aquatic Life use designation. During surveys in 2008 and 2010, Thomas (2008, p. 4) and USFWS (2010, pp. 1-13) observed Kentucky arrow darters in only one of these streams, Lower Buffalo Creek.

Middle Fork Kentucky River (3/3). This basin has fewer Kentucky arrow darter records than the North and South Forks of the Kentucky River. Much of the basin has been severely impacted by surface and underground coal mining, and an approximate 31-mile (50 km) segment of the Middle Fork has been influenced by the construction and inundation of Buckhorn Lake. During surveys from 2007 to 2009 (Thomas 2008, pp. 4-5; USFWS 2009, pp. 1-4), arrow darters were observed at two streams, Hell for Certain Creek (Leslie County) and Big Laurel Creek (Harlan County). Hell for Certain Creek lies at the western edge of the basin where the majority of its watershed is situated within the DBNF (Redbird District). Big Laurel Creek is a tributary of Greasy Creek at the southeastern edge of the basin (Thomas 2008, pp. 4-5; USFWS 2009, pp. 1-4). Despite being surrounded by what appeared to be an intact forest, habitat quality in Big

Laurel Creek was poor with heavy sedimentation (sand) and bank erosion (USFWS 2009, pp. 1-4). The source of the sediment was unknown, but upstream logging and associated road building were suspected as potential sources. Thomas (2008, p. 4) and USFWS (2009, pp. 1-4; 2010, pp. 1-13) surveyed 6 of 11 historic sites and 5 new sites in the Middle Fork Kentucky River basin and observed Kentucky arrow darters at only 2 of those sites. In March 2013, biologists with KDFWR and DBNF discovered an unknown population of *E. spilotum* in Laurel Creek, a second order tributary of Rockhouse Creek. Similar to Hell for Certain Creek, Laurel Creek is situated at the western edge of the Middle Fork basin and about 90 percent of its watershed is located within the DBNF (Red Bird Ranger District).

North Fork Kentucky River (12/22). The best habitat within this basin is located on UKs Robinson Forest in Breathitt and Knott Counties (Troublesome Creek basin). Two streams on Robinson Forest, Clemons Fork and Coles Fork, have stable populations of Kentucky arrow darters. The basins are intact and densely forested, with only minor interruption by logging roads. Both streams are high-gradient, cold, and dominated by cobble, boulder, and bedrock substrates. From 2007 to 2009, Thomas (2008, p. 5) and USFWS (2009, pp. 1-4) collected a total of 27 and 10 arrow darters, respectively, from Clemons Fork and Coles Fork. Thomas (2008, p. 5) and USFWS (2009, pp. 1-4) visited 12 additional Troublesome Creek basin sites located near Robinson Forest, but no arrow darters were collected. These sites were located on Bear Branch, Boughcamp Branch (of Buckhorn Creek), Buckhorn Creek, Balls Fork, Laurel Fork, Leatherwood Creek, Lewis Fork, Long Fork, Lost Creek, and Troublesome Creek. Kentucky arrow darters were first recorded from these basins during the 1960s, but these streams have been impacted severely by water pollution associated with coal mining activities, and portions of at least three of these streams, Buckhorn Creek (mile 0-6.8), Long Fork (mile 0-8.95), and Troublesome Creek (mile 0-45.1), have been placed on Kentuckys 303d list of impaired waters (KDOW 2008, pp. 65-101).

In 2010, a new but isolated population of Kentucky arrow darters was discovered in the headwaters of Buckhorn Creek, just upstream of its confluence with Emory Branch in Knott County (USFWS 2010, pp. 1-13). As mentioned above, much of the Buckhorn Creek mainstem has been degraded due to inputs of dissolved solids, sulfates, and sediment associated with mining activities in the surrounding watersheds. Instream conductivity values within Buckhorn Creek and many of its tributaries are elevated (typically exceeding 2,000 μS), and instream habitats have been degraded by sedimentation. During field surveys in August 2010, we observed similar water quality and habitat conditions downstream of Emory Branch (i.e., instream conductivity exceeded 2,000 μS , sedimentation was evident), but conditions upstream of Emory Branch were better, with low instream conductivity (110 μS) and suitable instream habitat. We observed three Kentucky arrow darters within this reach, which extends downstream from Prince Fork approximately 450 m (1,500 ft) to Emory Branch. Upstream of this reach, Buckhorn Creek is too shallow to support arrow darters.

Outside of the Troublesome Creek basin, Kentucky arrow darters were observed at two of eight sites in the Quicksand Creek basin (Breathitt and Knott Counties), four of six sites in the Frozen Creek basin, three of five sites in smaller tributaries of the lower North Fork Kentucky River basin, and one direct tributary (Silver Creek) of the Kentucky River just downstream of its confluence with the South Fork Kentucky River. Only one of these streams, Frozen Creek (Breathitt County), appeared to have good, stable habitat conditions (Thomas 2008, p. 5). The remaining streams with arrow darters were moderately to severely impacted by sedimentation, bank erosion, and riparian disturbance. Portions of two streams, Cope Fork and Hell Creek, have been identified as impaired by the Kentucky Division of Water and are listed on Kentuckys 303d list of impaired waters (KDOW 2008, pp. 65-101).

Population Estimates/Status:

Population estimates for the Kentucky arrow darter are not available; however, recent survey data (Thomas 2008, pp. 3-6; USFWS 2009, pp. 1-4) revealed that sites with arrow darters had an average of only three individuals per 100-m (328-ft) sampling reach and a median of two individuals per reach (range of 1 to 10 individuals). Recent surveys by the USFS (Brandt, pers. comm., 2011) from Laurel Fork and Cortland Branch of Left Fork Buffalo Creek (South Fork Kentucky River basin) produced slightly higher capture rates

(an average of 5 darters per 100-m (328-ft) sampling reach). The low abundance values (compared to other darters) are not surprising since both arrow darter species generally are not observed in large numbers, even in those streams where disturbance has been minimal (M. Thomas, pers. comm., 2010). The largest populations of Kentucky arrow darters were located in the following streams/basins:

Several tributaries of South Fork Kentucky River, Redbird District of DBNF (Clay and Leslie Counties); Hell Creek, Walker Creek, and Frozen Creek - direct tributaries of North Fork Kentucky River (Breathitt and Lee Counties);

Clemons Fork and Coles Fork of Buckhorn Creek, North Fork Kentucky River basin (Breathitt County).

Distinct Population Segment(DPS):

N/A

Threats

A. The present or threatened destruction, modification, or curtailment of its habitat or range:

The overall decline of the Kentucky arrow darter can be attributed to a variety of human-related activities in the upper Kentucky River basin. Activities such as coal mining, silviculture, agriculture, gas/oil well exploration, human development, and inadequate sewage treatment have all contributed to the degradation of streams within the range of the species (Branson and Batch 1972, pp. 513-516; Branson and Batch 1974, pp. 82-83; KDOW 2008, pp. 65-101; Thomas 2008, pp. 6-7). Adverse impacts result primarily from inputs of dissolved solids and elevation of instream conductivity, sedimentation, removal of riparian vegetation, bank erosion and channel instability, inputs of untreated sewage, and agricultural runoff.

Coal Mining

Coal mining activities represent the most imminent and substantial source of threats to the subspecies because these activities have the potential to significantly, and often permanently, alter instream water quality and cause physical habitat disturbance. Numerous studies have documented the fact that streams receiving discharge from mined areas exhibit characteristics not observed in unmined watersheds: (1) altered water quality conditions (Curtis 1973, pp. 153-155; Dyer and Curtis 1977, pp. 10-13; Dyer 1982, pp. 1-16; Hren et al. 1984, pp. 5-34; US EPA 2003, pp. 77-84; Pond et al. 2008, pp. 721-723); (2) increased sediment loads (Branson and Batch 1972, p. 513; Parker and Carey 1980, pp. 33-49 ; Osterkamp et al. 1984, pp. 59, 63; Pond 2004, pp. 19-20); (3) increased hydrologic response time to storm events (Bryan and Hewlett 1981, p. 298); (4) altered flow duration curves (USGS 2001, pp. 16-17); and (5) altered or changed channel morphology. As of March 2010, over 465 mining permits were active in the upper Kentucky River basin (LaSage, pers. comm., 2010). Some of these permits were active and coal removal was still occurring (about 360 permits), while others were inactive with reclamation activities underway (about 106 permits).

Impacts to instream water quality (chemistry) occur through inputs of dissolved metals and other solids that elevate stream conductivity, increase sulfate levels, and/or increase stream pH, (Curtis 1973, pp. 153-155; Pond 2004, pp. 6-7, 38-41; Hartman et al. 2005, p. 95; Mattingly et al. 2005, p. 59; Palmer et al. 2010, pp. 148-149). As rock strata and overburden (excess material) are exposed to the atmosphere, precipitation leaches metals and other solids (e.g., Calcium, Magnesium, Sulfates, Iron, Manganese) from these materials and carries them in solution to receiving streams (Pond 2004, p 7). If valley fills are used as part of the mining activity, precipitation and groundwater percolate through the fill and dissolve minerals until they discharge at the toe of the fill as surface water (Pond et al. 2008, p. 718). Both of these scenarios result in elevated conductivity, sulfates, and hardness in the receiving stream. Increased levels of these metals and

other dissolved solids have been shown to exclude fish species from streams in eastern Kentucky, including the federally threatened blackside dace (*Chrosomus cumberlandensis*) in the upper Cumberland River basin (Mattingly et al. 2005, pp. 59-62).

Based on earlier research by Branson and Batch (1974, pp. 81-83) and Dyer and Curtis (1977, pp. 1-13) and recent fish survey results by Thomas (2008, pp. 3-6) and USFWS (2009, pp. 1-4), it is clear that degraded water quality conditions in the upper Kentucky River basin have adversely affected Kentucky arrow darter populations. From late 1967 to 1975, Branson and Batch (1972, pp. 507-518; 1974, pp. 81-83), and Dyer and Curtis (1977, pp. 1-13) studied the effects of strip mining activities on water quality and stream fishes in the Quicksand Creek (Leatherwood Creek) and Buckhorn Creek (Bear Branch) basins, Breathitt County. Six first-order watersheds, three in the Leatherwood Creek basin and three in the Bear Branch basin, were investigated during the study, beginning in late summer 1967 prior to the onset of mining and continuing until 1975. One of the six small watersheds, Jenny Fork, was not mined and served as a control watershed. Water quality data from mined watersheds showed increases in conductivity, sulfate, magnesium, bicarbonate, and silt deposition (Dyer and Curtis 1977, pp. 3-7, 13). Water quality data from the reference site, Jenny Branch, showed little variation and remained at baseline levels.

Fish community data from the Bear Branch and Leatherwood Creek watersheds showed that fishes were pushed downstream or eliminated from the fauna altogether in mined watersheds (Branson and Batch 1972, pp. 514-515; Branson and Batch 1974, pp. 82-83). The only exception to this was the creek chub, which appeared to be tolerant of mining impacts. Several species, silver shiner (*Notropis photogenis*), Kentucky arrow darter, Johnny darter, variegated darter (*Etheostoma variatum*), greenside darter (*E. blennioides*), and emerald darter were eliminated from Leatherwood Creek. Two species, northern hogsucker (*Hypentelium nigricans*) and blackside darter (*Percina maculata*), were eliminated from both streams. During the last fish sampling event in September 1972, Kentucky arrow darters were observed at the mouth of Bear Branch (Branson and Batch 1974, p. 82), but instream conductivity levels had not peaked. Branson and Batch (1972, p. 514) also did not observe young darters and minnows during later visits (early 1970s), suggesting that reproduction had been curtailed by the mining activity. Thomas (2008, p. 5) and USFWS (2009, pp. 1-4) resurveyed these streams in 2008 and found that conductivity levels had increased since the 1970s, reaching 845 μS in Bear Branch and 1008 μS in Leatherwood Creek. Kentucky arrow darters were not observed at these sites.

Recent range-wide surveys by Thomas (2008, pp. 3-6) and USFWS (2009, pp. 1-4) demonstrated that Kentucky arrow darters are excluded from watersheds when conductivity levels exceed about 250 μS . Mattingly et al. (2005, pp. 59-62) reported virtually identical results for the federally threatened blackside dace in the upper Cumberland River basin. Historic arrow darter sites that lacked darters had higher conductivity values (average = 680 μS) than historic sites that continued to support arrow darters (average = 105 μS). Arrow darters were observed at only one historic site (USFWS 2009, pp. 1-4), Walker Creek (Owsley County), with a conductivity value greater than 250 μS (400 μS).

There is a pattern of increasing conductivity and loss of arrow darter populations which is evident in the fish and water quality data from the Buckhorn Creek basin (1962 to present) in Breathitt and Knott Counties. Kentucky arrow darters and other fish species were first reported from the basin in 1962 by Kuehne (1962, p. 608-609), who surveyed sites on the Buckhorn Creek mainstem and numerous tributaries - Bear Branch, Clemons Fork, Coles Fork, Laurel Fork, Lewis Fork, and Long Fork. Kuehne (1962, p. 608-609) documented Kentucky arrow darters at 16 of 22 sites within the basin. Since that time, the majority of these watersheds have been mined extensively and conductivities have increased, especially in areas to the south and east of the Buckhorn Creek mainstem. The only exceptions are two unmined watersheds on UK's Robinson Forest (Clemons Fork and Coles Fork) and an approximate 450-m (1,500-ft) reach in the headwaters of Buckhorn Creek. Thomas (2008, p. 5) and USFWS (2009, pp. 1-4; 2010, pp. 1-13) resurveyed sites on all historic streams (and most historic sites) in the Buckhorn Creek basin from 2007 to 2010, observing Kentucky arrow darters in only Clemons Fork, Coles Fork, and Buckhorn Creek, upstream of Emory Branch. Conductivity levels of Clemons Fork, Coles Fork, and Buckhorn Creek (upstream of Emory Branch) remained at or near

background levels, (50 to 110 μS), but conductivity levels at other streams were elevated, with some of these being exceptionally high. Portions of two of these streams, Buckhorn Creek (mile 0-6.8) and Long Fork (mile 0-8.95) have been placed on Kentucky's 303d list of impaired waters (KDOW 2008, pp. 65-101).

Mine drainage also causes physical and chemical impacts to streams as a result of the precipitation of entrained metals and sulfate, which become unstable in solution (USEPA 2003, pp. 77-84; Pond 2004, p. 7). Hydroxide precipitants are formed from iron and aluminum, creating orange or white sludge (yellow boy) that forms a thick coating on stream substrates (Pond 2004, p. 7). Most affected streams also have elevated levels of calcium in solution, and if pH is elevated, calcium sulfate (CaSO_4) or calcium carbonate (CaCO_3) will precipitate (USEPA 2003, pp. 77-84; Pond 2004, p. 7). These precipitants accumulate on substrates, encrusting and cementing stream sediments, making them unsuitable for colonization by invertebrates and rendering them unsuitable as foraging or spawning habitat for the Kentucky arrow darter. Acid mine drainage (AMD) tends to be more of a legacy problem, as enforcement, newer technology, and mining methods have mostly eliminated it in the coal fields of Kentucky and Tennessee (Pond 2004, p. 6). In the few streams where the problem persists, AMD can be highly detrimental to fish and aquatic insect populations (Henry et al. 1999, pp. 919-920; Pond 2004, pp. 7-8). Streams affected by AMD tend to have low pH, high conductivity, and high metal and sulfate concentrations (Herlihy et al. 1990, pp. 101-105; Pond 2004, pp. 7-8).

Silviculture

Logging activities can adversely affect Kentucky arrow darters through sedimentation of instream habitats, removal of streamside (riparian) vegetation, and direct channel disturbance. Sedimentation occurs as soils are disturbed, the overlying leaf or litter layer is removed, and sediment is carried overland from logging roads, stream crossings, skid trails, and riparian zones during storm events. Excess sediment can bury instream habitats used by the species for foraging, reproduction, and sheltering, and it can disrupt the dynamic equilibrium of channel width, depth, flow velocity, discharge, channel slope, roughness, sediment load, and sediment size that maintains stable channel morphology. This can lead to channel instability and further degradation of instream habitats. Reductions in riparian vegetation can adversely affect the species through increased solar radiation, elevated stream temperatures, loss of allochthonous (organic material originating from outside the channel) food material, and bank instability / erosion. Direct channel disturbance occurs primarily at stream crossings during culvert, log, or rock placement. Severe impacts can occur when loggers use stream channels illegally as skid trails (M. Floyd, pers. obs. 2009).

Sediment (Nonpoint-Source)

Sediment (siltation) has been listed repeatedly by the Kentucky Natural Resources and Environmental Protection Cabinet (Division of Water) as the most common stressor of aquatic communities in the upper Kentucky River basin (KDOW 2008, pp. 65-101). Sedimentation comes from a variety of sources, but KDOW identified the primary sources of sediment as loss of riparian habitat, surface coal mining, and legacy coal extraction (KDOW 2008, pp. 65-101). All of these activities can result in canopy removal, channel disturbance, and increased siltation, thereby degrading habitats used by fishes for both feeding and reproduction. The reduction or loss of riparian vegetation results in the elevation of stream temperatures, destabilization of stream banks and siltation, and removal of submerged root systems that provide habitat for fish and macroinvertebrates (Mattingly et al. 2005, p. 5). Unpaved (dirt, gravel) roads represent another sediment source that can deliver large amounts of silt or sand during storm events. Many of these roads are poorly ditched, which creates eroded channels within the roadway that can carry sediment directly to nearby streams. Numerous streams within the Kentucky arrow darters current range have been identified as impaired (primarily due to siltation from mining, logging, agricultural activities, and land development) and have been included on Kentucky's 303(d) list of impaired waters (KDOW 2008, pp. 65-101).

Sediment has been shown to damage and suffocate fish gills and eggs, larval fishes, bottom dwelling algae, and other organisms; reduce aquatic insect diversity and abundance; and, ultimately, negatively impact fish growth, survival, and reproduction (Waters 1995, pp. 5-7; Meyer and Sutherland (2005, pp. 2-3). Wood and Armitage (1997, pp. 211-212) identified at least five impacts of sedimentation on fish, including (1)

reduction of growth rate, disease tolerance, and gill function; (2) reduction of spawning habitat and egg, larvae, and juvenile development; (3) modification of migration patterns; (4) reduction of food availability through the blockage of primary production; and (5) reduction of foraging efficiency. Kentucky arrow darter habitats are also affected when riparian corridors are disturbed or significantly altered during mine preparation, logging activities, or road construction.

Oil and Gas

Oil and gas exploration activities represent a threat to the species. Exploration and drilling activities typically involve the construction of new roads and stream crossings, so the potential is high for sedimentation and other direct, physical disturbance to stream channels (e.g., culvert placement, road building). Water quality degradation is a more significant threat associated with these activities because releases of chemicals can occur during the drilling process. Significant releases from gas well sites have been documented within the upper Cumberland River basin (M. Floyd, pers. obs., 2009) and likely occur in the Kentucky River basin as well.

Walker Creek, a direct tributary of the North Fork Kentucky River, is the only stream within the subspecies range that continues to support arrow darters despite having a conductivity value (400 μS) greater than 250 μS . At the time of the field survey, the source of the elevated conductivity was unknown, but subsequent review of oil and gas mapping revealed that legacy oil and gas wells were abundant in the watershed, and along with legacy mining impacts, likely have contributed to the elevated conductivity. An adjacent stream, Hell Creek, has been placed on Kentucky's 303d list due to impacts from abandoned mines and petroleum/gas production activities.

Other

Other nonpoint-source pollutants that affect the Kentucky arrow darter include domestic sewage (through septic tank leakage or straight pipe discharges) and agricultural pollutants such as animal waste, fertilizers, pesticides, and herbicides. Nonpoint-source pollutants can cause excess eutrophication (increased levels of nitrogen and phosphorus) (Table 3, Laurel Creek), excessive algal growth, instream oxygen deficiencies, and other changes in water chemistry that can seriously impact aquatic species (KDOW 1996, pp. 48-50; KDOW 2006, pp. 70-73).

B. Overutilization for commercial, recreational, scientific, or educational purposes:

The Kentucky arrow darter is not believed to be utilized for commercial, recreational, scientific, or educational purposes. Individuals may be collected occasionally in minnow traps and used as live bait, but this activity does not pose a significant threat.

C. Disease or predation:

Although the Kentucky arrow darter is undoubtedly consumed by predators, predation by naturally occurring predators is a normal aspect of the population dynamics and is not considered to currently pose a threat to the species.

D. The inadequacy of existing regulatory mechanisms:

The Commonwealth of Kentucky prohibits the collection of the Kentucky arrow darter and other fish species for scientific purposes without a valid state-issued collecting permit. However, this requirement does not provide any protection to the species habitat. Within Kentucky, persons who hold a valid fishing license (obtained from Kentucky Department of Fish and Wildlife Resources (KDFWR)) are allowed to collect up to 500 minnows per day (a minnow is defined as any non-game fish less than 6 inches in length, with the exception of federally listed species). This regulation allows for the capture, holding, and potential use of the Kentucky arrow darter as a bait species. While we do not currently believe this is a significant threat (see Factor B), it is a potential threat.

The Kentucky arrow darter and its habitats are afforded some protection from water quality and habitat degradation under the Clean Water Act of 1977 (33 U.S.C. 1251 et seq.), Surface Mining Control and Reclamation Act of 1977 (30 U.S.C. 1234-1328), Kentucky's Forest Conservation Act of 1998 (KRS 149.330-355), Kentucky's Agriculture Water Quality Act of 1994 (KRS 224.71-140), and additional Kentucky laws and regulations regarding natural resources and environmental protection (KRS 146.200-360; KRS 224; 401 KAR 5:026, 5:031). However, as demonstrated under Factor A, population declines and degradation of habitat for this species are ongoing despite the protection afforded by these laws and corresponding regulations. While these laws have resulted in some improvements in water quality and stream habitat for aquatic life, including the Kentucky arrow darter, they alone have not been adequate to fully protect this species; elevated conductivity, sedimentation, and non-point source pollutants continue to be a problem.

States maintain water-use classifications through issuance of National Pollutant Discharge Elimination System (NPDES) permits to industries, municipalities, and others that set maximum limits on certain pollutants or pollutant parameters. For water bodies on the 303(d) list, States are required under the Clean Water Act to establish a total maximum daily load (TMDL) for the pollutants of concern that will bring water quality into the applicable standard. Eighteen (18) Kentucky arrow darter streams have been identified as impaired by the Kentucky Division of Water and placed on the State's 303(d) list (KDOW 2008) (Table 3). Causes of impairment were listed as increased sediment/siltation and total dissolved solids from coal mining, silviculture, loss of riparian habitat, and agriculture, and organic enrichment/eutrophication from agriculture. TMDLs have not yet been developed for these pollutants.

The Kentucky arrow darter has been designated as Threatened in Kentucky by the KSNPC (KSNPC 2005), but this designation conveys no legal protection under Kentucky state law.

E. Other natural or manmade factors affecting its continued existence:

The disjunct nature of some Kentucky arrow darter populations prohibits the natural interchange of genetic material between populations, and the small population size reduces the reservoir of genetic diversity within populations. This can lead to inbreeding depression and reduced fitness of individuals (Soule 1980, pp. 157-158; Hunter 2002, pp. 97-101). It is possible that some of the arrow darter populations are below the effective population size required to maintain long-term genetic and population viability (Soule 1980, p. 162-164; Hunter 2002, pp. 105-107).

Climate change has the potential to increase the vulnerability of the Kentucky arrow darter to random detrimental events (e.g., McLaughlin et al. 2002; Thomas et al. 2004). Global warming is expected to result in increasing frequency and duration of droughts and the strength of storms (e.g., Cook et al. 2004). Severe droughts similar to those that affected eastern Kentucky in 2007 and 2008 could be intensified by rising mean air temperatures and reduced precipitation amounts as predicted by Mauer et al. (2007) and ClimateWizard (2009) over the next 40 years in eastern Kentucky.

Conservation Measures Planned or Implemented :

Propagation/Reintroduction Effort Clay and Leslie Counties, DBNE. KDFWR identified the Kentucky arrow darter as a Species of Greatest Conservation Need (SGCN) in its State Wildlife Action Plan (KDFWR 2005, p. 2.2.2). The plan identifies conservation issues (threats), conservation actions, and monitoring strategies for 251 animal species belonging to one of 20 terrestrial and aquatic habitat guilds (collection of species that occur in the same habitat). To fully understand these conservation issues, the KDFWR developed a priority list of research and survey needs for Kentucky's SGCN. The KDFWR attempted to address two of these needs in 2008 by initiating a propagation and reintroduction study for the Kentucky arrow darter through the Services State Wildlife Grant program (Ruble et al. 2010, pp. 1-8). The study was designed to document details on the subspecies reproductive biology and to begin conservation actions (e.g., propagation and

augmentation) that would preclude the need to list the Kentucky arrow darter as threatened or endangered under the ESA. The KDFWR partnered with Conservation Fisheries, Inc. (CFI) to develop successful spawning protocols and produce the offspring needed to augment populations within the subspecies current range.

From 2009 to 2011, a total of 145 captive-spawned, juvenile arrow darters (from Big Double Creek) were produced by CFI and released into Sugar Creek, Leslie County, a tributary of the Red Bird River in the DBNF, Redbird District (Thomas and Brandt 2012, pp. 1-11). Sugar Creek was chosen as a release site because it was located on the DBNF, it was within the subspecies known range, and it contained suitable habitat. Prior to release, all arrow darters were marked at CFI with a Northwest Marine Technologies elastomer tag. Attempts to relocate tagged darters were made in August 2009, October 2009, March 2010, January 2012, and February 2012. Unfortunately, no tagged darters were observed. A total of 31 untagged arrow darters were observed in Sugar Creek (5 in 2009, 4 in 2010, and 22 in 2012), so KDFWR and CFI made the decision to abandon efforts at Sugar Creek and begin another reintroduction effort at Long Fork, another DBNF stream and tributary of Hector Branch in Clay County.

In August and October 2012, a total of 829 tagged Kentucky arrow darters (about 50-55 mm total length) were introduced within a 1.5-km reach of Long Fork. Monitoring has been conducted on two occasions since the initial release using visual searches and seining methods. The initial monitoring effort (October 2012) produced a total of 18 tagged darters. The second effort (January 2013) was more successful, producing a total of 47 tagged darters. Most of these individuals were found in pools (depth of 8-24 inches) with rock substrates, exposed bedrock, and some marginal cover (e.g., tree roots). During the January 2013 survey, no individuals were captured downstream of a waterfall barrier (about 100 m above the mouth), but tagged darters began to appear about 100 to 200 m upstream of the waterfall, and capture rates began to increase as the surveyors moved upstream. About 2-3 individuals were observed per pool below the mid-point of the reach, increasing to 4-6 per pool above the midpoint. KDFWR and CFI completed another monitoring event on March 13, 2013, when about 50 tagged darters were observed. All fish were in very good condition and had grown in size. Surveys were concentrated upstream of the midway point and fish were captured up to about 200 m upstream of the most upstream release point. Additional monitoring and releases are planned for the summer/fall of 2013.

Candidate Conservation Agreement (CCA) with DBNE. The KFO and the DBNF are working cooperatively to develop a candidate conservation agreement for the Kentucky arrow darter on the DBNF. Over half of the species extant streams occur on lands owned and managed by the DBNF, so conservation of these populations is essential to the species recovery, and a DBNF-specific conservation plan is needed to guide those efforts. The DBNF and KFO met in January 2013 to discuss potential conservation actions, including (1) an evaluation of relevant Forest Plan standards to identify any impediments to KAD conservation and ways to address those impediments, (2) a review of available data that would help identify any immediate opportunities for conservation success with the KAD, (3) the development of a watershed-based threat matrix for USFS lands and watersheds, (4) identifying and ranking potential stream restoration opportunities, (5) evaluating the implications of rainbow trout stocking within USFS lands and watersheds, and (6) the development of a Candidate Conservation Agreement for the DBNF. Both agencies are currently working on action items, and completion of a draft CCA is scheduled for July 2013.

Range-wide Candidate Conservation Agreement with Assurances (CCAA). The KFO is assisting the KDFWR in development of a range-wide CCAA for the species. To date, the process has included a Science Advisory Committee (SAC) meeting in January 2013 and a Stakeholder meeting in February 2013. The SAC meeting was moderated by KDFWR and included a total of 18 biologists, representing a variety of agencies/groups: Appalachian Wildlife Foundation, Appalachian Technical Services, Austin Peay State University, Conservation Fisheries Inc., DBNF, KDFWR, Kentucky Division of Water, KSNPC, Morehead State University, The Nature Conservancy, U. S. Environmental Protection Agency, USFWS (KFO), and the

U. S. Geological Survey. The SAC reviewed and discussed the species biology and current status/threats and identified three general research needs or priorities for the species: (1) genetics, (2) stressor identification, and (3) dispersal behavior (movement).

The Stakeholder meeting was led by staff from KDFWR and was attended by a variety of stakeholders from the coal mining, forestry, transportation, and environmental consulting industries, the DBNF, the Appalachian Wildlife Foundation, Eastern Kentucky University, and KSNPC. Dr. Michael Floyd (KFO) presented information on KAD life history and threats and gave a summary of CCAAs and CCAA development. The meeting also included a discussion session where a variety of questions by potential stakeholders were answered by KDFWR and/or KFO staff. The KFO and KDFWR continue to work on development of the CCAA; the KFO is currently working on a Draft Range-Wide Conservation Strategy for the species.

Upper Kentucky River Basin Distributional Study and Habitat Characterization. The KFO is working cooperatively with the KSNPC to investigate the distribution, status, population size, and environmental resource use of the Kentucky arrow darter within the upper Kentucky River system. One important aspect in assessing these components is to account for imperfect detection when surveying for the species. Studies that do not account for imperfect detection can often lead to an underestimation of the true proportion of sites occupied and can bias assessments and efforts (MacKenzie et al. 2002, pp. 2248-2255; MacKenzie et al. 2004, pp. 149-172). The KSNPC and KFO propose to address these concerns by meeting the following objectives:

- Estimate occupancy and detection probability of the focal species;
- Estimate the proportion of sites occupied by the focal species, after accounting for the detection probability;
- Draw inference into the environmental resources important to the persistence of the focal species;
- Assess the status of the focal species;
- Estimate the population size of the species within the study area; and
- Make recommendations for future efforts in the conservation of the focal species.

Eastern Kentucky University (EKU) and KFO Movement Study and Population Estimate. The KFO is working with EKU (Dr. Sherry Harrel and graduate student, Mr. Jonathan Baxter) to develop and implement a movement study, habitat characterization, and population estimate of two Kentucky arrow darter streams, Gilberts Big Creek and Elisha Creek (DBNF), in Clay and Leslie Counties (Harrel and Baxter 2012, pp. 1-12). Mr. Baxter will use PIT-tags and placed antenna systems to monitor intra- and inter-tributary movement patterns in both streams, and he will be collecting seasonal (spring, summer, and fall) biotic and abiotic data from 17 100-m reaches to determine habitat use and population density/size for both streams.

Landscape Model of KAD Occurrence in Response to Water Quality. Through USFWS-USGS Quick Response funding, the Kearneysville, West Virginia USGS office and KFO are working cooperatively to apply spatial statistical modeling techniques to estimate Kentucky arrow darter distribution and sensitivity to water quality parameters (Hitt and Floyd 2012, pp. 1-7). The study has three specific objectives: (1) Evaluate stream conductivity relations to land use in the study area using spatial analysis and modeling techniques; (2) Assess potential threshold responses of Kentucky arrow darter occurrence to conductivity and other water quality parameters; and (3) Develop and test spatially explicit species distribution models for Kentucky arrow darter using an integrated fish and water quality dataset from KDFWR and USFWS.

Bullskin Creek Restoration, Leslie County. A 0.55-km (1823-ft) stream restoration/enhancement project was completed in 2005 in the upper reaches of Bullskin Creek, Leslie County. Bullskin Creek represents suitable habitat for the Kentucky arrow darter and the subspecies was reported from the Bullskin Creek watershed as recently as 2007, when 2 individuals were collected approximately 12.1 stream km [7.5 mi] downstream of the restoration site (Thomas 2008, p. 4). The Bullskin project was funded through Kentuckys Wetland and Stream Mitigation Fund (managed by KDFWRs Stream and Wetland Restoration Program) and was intended to repair eroding banks and poor habitat conditions within Bullskin Creek. The project included a permanent,

9-m (30-foot) easement held by KDFWR. Habitat improvements in this reach of Bullskin Creek will benefit Kentucky arrow darters living in upstream and downstream reaches.

Summary of Threats :

Three of the five listing factors considered by the Service pose threats to the Kentucky arrow darter: the present or threatened destruction, modification or curtailment of its habitat or range; the inadequacy of existing regulatory mechanisms; and other natural or manmade factors affecting its continued existence. The species habitat and range have been severely degraded and limited by water pollution from surface coal mining and gas exploration activities; removal of riparian vegetation; stream channelization; increased siltation associated with poor mining, logging, and agricultural practices; and deforestation of watersheds. These threats are (1) widespread across the subspecies range (the geographic scope is widespread and not localized); (2) imminent (the effects are manifested immediately and will continue); and (3) severe (stressors are of high intensity or high strength and can lead to mortality). The severity (or intensity) of these threats, especially impacts from mining and gas exploration activities, is high these activities can permanently alter stream water quality (e.g., elevated conductivity) by contributing sediment, dissolved metals, and other solids to streams supporting Kentucky arrow darter populations. These water quality changes can be permanent and render these habitats unsuitable for the Kentucky arrow darter. Recent and past research has demonstrated that the subspecies is intolerant of these conditions, and it has been eliminated from a number of streams across its range. Current regulatory mechanisms have been inadequate to prevent these impacts. The small, remnant nature of many Kentucky arrow darter populations may prohibit the natural interchange of genetic material between these populations, and the small population size may reduce the reservoir of genetic diversity within populations. This can lead to inbreeding depression and reduced fitness of individuals. It is possible that some Kentucky arrow darter populations are below the effective population size required to maintain long-term genetic and population viability. We have no information indicating that the magnitude or imminence of these threats is likely to be appreciably reduced in the foreseeable future.

We evaluated the threats to the Kentucky arrow darter and considered factors that, individually and in combination, presently or potentially could pose a risk to the subspecies and its habitat. Based on our analysis of these threats, we find that this species is warranted for listing throughout all its range, and, therefore, find that it is unnecessary to analyze whether it is threatened or endangered in a significant portion of its range.

For species that are being removed from candidate status:

_____ Is the removal based in whole or in part on one or more individual conservation efforts that you determined met the standards in the Policy for Evaluation of Conservation Efforts When Making Listing Decisions(PECE)?

Recommended Conservation Measures :

Conservation measures for the Kentucky arrow darter should be concentrated in those watersheds where populations still occur and should be focused primarily on the protection and restoration of the existing water and habitat quality of these systems. Seven conservation actions were developed for the subspecies habitat guild as part of KDFWRs State Wildlife Action Plan (KDFWR 2005, p. 2.2.2): (1) the creation of financial incentives to protect riparian corridors and watersheds, (2) acquisition and conservation easements of critical aquatic habitat, (3) encouragement and assistance in developing and implementing best management practices, (4) restoration of degraded habitats, (5) coordination and implementation of existing Farm Bill programs or other federal incentive programs, (6) education of user groups on significance and importance of riparian corridors and watersheds, and (7) development and initiation of local watershed improvement projects. Other appropriate conservation actions identified in KDFWRs plan include the development of protection and enhancement (mitigation) plans for mined watersheds and the development of strategies for reintroduction and enhancement of populations. In order to achieve conservation goals, the Service could pursue the development of candidate conservation agreements and candidate conservation agreements with

assurances with potential partners. Conservation efforts could be augmented through additional research on the species current distribution, life history, environmental requirements, and movement patterns.

Priority Table

Magnitude	Immediacy	Taxonomy	Priority
High	Imminent	Monotypic genus	1
		Species	2
		Subspecies/Population	3
	Non-imminent	Monotypic genus	4
		Species	5
		Subspecies/Population	6
Moderate to Low	Imminent	Monotype genus	7
		Species	8
		Subspecies/Population	9
	Non-Imminent	Monotype genus	10
		Species	11
		Subspecies/Population	12

Rationale for Change in Listing Priority Number:

Based on new morphological and genetic analyses by Thomas and Johansen (2008, p. 46) and published species accounts and lists in Page and Burr (2011, p. 569) and Eschmeyer (2013), the Kentucky arrow darter is now recognized as *E. spilotum* Gilbert. The elevation to species rank increases the listing priority from 3 (subspecies) to 2 (species).

Magnitude:

The severity (or intensity) of these threats, especially impacts from mining, is high. These activities can permanently alter stream water quality (e.g., elevated conductivity) by contributing sediment, dissolved metals, and other solids to streams supporting Kentucky arrow darter populations and these threats are widespread across the species' range (the geographic scope is widespread and not localized). These water quality changes can be permanent and render these habitats unsuitable for the species. Recent and past research has demonstrated that the species is intolerant of these conditions and it has been eliminated from a number of streams across its range. Collectively, these factors are serious and significant impediments to the survival of the Kentucky arrow darter and are of "High" magnitude.

Imminence :

Threats to the Kentucky arrow darter are imminent because the effects are ongoing and will continue. Demand for coal production is not expected to decline over the next 10 years and may increase; consequently, water quality degradation and physical habitat disturbance associated with coal mining activities are expected to continue. Other threats to the subspecies are not expected to diminish in the foreseeable future. Therefore, these factors are "Imminent."

Yes No Have you promptly reviewed all of the information received regarding the species for the purpose

of determination whether emergency listing is needed?

Emergency Listing Review

 No Is Emergency Listing Warranted?

During the normal listing process, the subspecies will continue to be threatened by the listing factors described above, but these threats are not expected to jeopardize the subspecies' continued existence. The species' continued existence is bolstered by the fact that populations occurring on public land (DBNF, Red Bird District and Robinson Forest) are protected from severe habitat degradation associated with coal mining and logging activities.

Description of Monitoring:

The KDFWR (Thomas 2008, pp. 3-6) and the KFO (USFWS 2009, pp. 1-4) completed a basin-wide survey for the Kentucky arrow darter during the summers of 2007 to 2009. All streams known to support the species were visited, as well as 85 percent of the species historic sites. In 2010, surveys were initiated by the KFO within the Kentucky arrow darters historic range, including 1 historic stream (Buckhorn Creek) and 14 other streams lacking previous records for the species (USFWS 2010, pp. 1-13). An isolated population of Kentucky arrow darters was discovered in the headwaters of Buckhorn Creek, but no Kentucky arrow darters were observed at the 14 new streams. In April 2013, KDFWR and DBNF biologists discovered a new population of Kentucky arrow darters in Laurel Creek, a second order tributary of Rockhouse Creek in Leslie County (Thomas pers. comm. 2013). Laurel Creek lies at the western edge of the Middle Fork basin, and about 90 percent of its watershed is located within the DBNF (Red Bird Ranger District).

Indicate which State(s) (within the range of the species) provided information or comments on the species or latest species assessment:

Kentucky

Indicate which State(s) did not provide any information or comment:

none

State Coordination:

Kentucky (Kentucky Department of Fish and Wildlife Resources and Kentucky State Nature Preserves Commission). The Kentucky arrow darter is included as a species of conservation concern in Kentucky's State Wildlife Action Plan (KDFWR 2005, p. 2.2.2).

Literature Cited:

Awise, J.C. and J.L. Hambrick, eds. 1996. Conservation genetics: case histories from nature. Chapman and Hall, New York.

Bailey, R. M. 1948. Status, relationships, and characters of the percid fish, *Poeciliichthys sagitta* Jordan and Swain. *Copeia* 1948:77-85.

Bailey, R. M. and W. A. Gosline. 1955. Variation and systematic significance of vertebral counts in the American fishes of the family Percidae. *Miscellaneous Publications, Museum of Zoology, University of Michigan* 93:1-44.

Bailey, R. M., H. E. Winn, and C. L. Smith. 1954. Fishes from the Escambia River, Alabama and Florida,

with ecologic and taxonomic notes. Proceedings of the Academy of Natural Sciences of Philadelphia 106:109-164.

Branson, B. A. and D. L. Batch. 1972. Effects of strip mining on small stream fishes in east-central Kentucky. Proceedings of the Biological Society of Washington. 84:507-518.

Branson, B. A. and D. L. Batch. 1974. Additional observations on the effects of strip mining on small-stream fishes in east-central Kentucky. Trans. Kentucky Acad. Sci. 35:81-83.

Branson, B. A. and D. L. Batch. 1983. Fishes of the South Fork of the Kentucky River, with notes and records from other parts of the drainage. Southeastern Fishes Council Proceedings 4:1-15.

Bryan, B.A., and Hewlett, J.D. 1981. Effects of surface mining on storm flow and peak flow from six small basins in eastern Kentucky: Water Resources Bulletin 17:290-299.

Burr, B. M. and M. L. Warren, Jr. 1986. A distributional atlas of Kentucky fishes. Kentucky State Nature Preserves Commission Scientific and Technical Series Vol. 4.

ClimateWizard. 2009. Models of average temperature and precipitation changes for Kentucky over the next 40 years. <http://www.climatewizard.org/>.

Cook, E.R., C.A. Woodhouse, C.M. Eakin, D.M. Meko, and D.W. Stahle. 2004. Long-term aridity changes in the western United States. Science 306:1015-1018.

Curtis, W. R. 1973. Effects of strip mining on the hydrology of small mountain watersheds in Appalachia. Pages 145-157 In R. J. Hutnik and G. Davis. Eds. Ecology and reclamation of devastated land. Vol. 1. Gordon and Breach, New York.

Dyer, K. L. and W. R. Curtis. 1977. Effect of strip mining on water quality in small streams in eastern Kentucky, 1967-1975. USDA Forest Service Research Paper NE-372. Northeastern Forest Experiment Station, Upper Darby, Pennsylvania. 13 pp.

Eschmeyer, W. N. (ed). 2012. Catalog of Fishes. California Academy of Sciences (<http://research.calacademy.org/research/ichthyology/catalog/fishcatmain.asp>). Electronic version accessed April 12, 2013.

Etnier, D.A. and W.C. Starnes. 1993. The Fishes of Tennessee. The University of Tennessee Press, Knoxville, Tennessee. 681 pp.

Evans, R. 2010. Personal communication regarding the status of the Kentucky arrow darter. Biologist, Kentucky State Nature Preserves Commission, Frankfort, Kentucky.

Gilbert, C. H. 1887. Descriptions of new and little known etheostomoids. Proceedings of the United States National Museum 10:47-64.

Greenberg, L. A. and S. Steigerwald. 1981. New distributional record for *Etheostoma sagitta* in Kentucky. Transactions of the Kentucky Academy of Science. 42:37.

Harrel, S. L. and J. B. Baxter. 2012. Distribution, movement, and ecology, of *Etheostoma sagitta spilatum* (Gilbert), the Kentucky arrow darter, in second and third order streams in two subbasins of the Kentucky River basin. Draft research proposal submitted to the U.S. Fish and Wildlife Service, Frankfort, Kentucky. 12 pp.

Hartman, K.J., M.D. Kaller, J.W. Howell, and J.A. Sweka. 2005. How much do valley fills influence

headwater streams? *Hydrobiologia* 532:91-102.

Henry, T. B., E. R. Irwin, J. M. Grizzle, M. L. Wildhaber, and W. G. Brumbaugh. 1999. Acute toxicity of an acid mine drainage mixing zone to juvenile bluegill and largemouth bass. *Transactions of the American Fisheries Society* 128:919-928.

Herlihy, A. T., P. R. Kaufmann, and M. E. Mitch. 1990. Regional estimates of acid mine drainage impact on streams in the Mid-Atlantic and southeastern United States. *Water, Air, and Soil Pollution* 50:91-107.

Hitt, N. P. and M. A. Floyd. 2012. Landscape modeling of Kentucky arrow darter (*Etheostoma sagitta spilotum*) occurrence and response to water quality parameters. USFWS and USGS Quick Response Program Proposal. Kearneysville, West Virginia and Frankfort, Kentucky. 7 pp.

Hren, J., K. S. Wilson, and D. R. Helsel. 1984. A statistical approach to evaluate the relation of coal mining, land reclamation, and surface water quality in Ohio. U.S. Geological Survey Water Resources Investigations Report 84-4117. 36 pp.

Hunter, M. L., Jr. 2002. *Fundamentals of conservation biology*, second edition. Blackwell Science, Inc. Malden, Massachusetts. 547 pp.

Jordan, D.S. and J. Swain. 1883. List of fishes collected in the Clear Fork of the Cumberland, Whitley County, Kentucky, with descriptions of three new species. *Proceedings of the United States National Museum* 6:248-251.

Kentucky Department of Fish and Wildlife Resources (KDFWR). 2005. *Kentucky's Comprehensive Wildlife Conservation Strategy*. KDFWR, Frankfort, Kentucky. <http://fw.ky.gov/kfwis/stwg/>.

Kentucky Division of Water (KDOW). 1996. Kentucky 305(b) report to Congress on water quality. Natural Resources and Environmental Protection Cabinet, Frankfort, Kentucky. 199 pp.

Kentucky Division of Water. 2006. *Integrated report to congress on water quality in Kentucky. Volume I. 305(b) assessment results with emphasis on the Kentucky River basin management unit and Salt-Licking Rivers basin management unit*. Natural Resources and Environmental Protection Cabinet, Frankfort, Kentucky. 222 pp.

Kentucky Division of Water. 2008. *Draft 2008 integrated report to Congress on the condition of water resources in Kentucky. Vol. II. 303(d) list of surface waters*. Kentucky Environmental and Public Protection Cabinet, Frankfort, Kentucky. 609 pp.

Kentucky State Nature Preserves Commission. 2005. *Rare and extirpated biota of Kentucky*. (pdf file available at: www.naturepreserves.ky.gov). 19 pp.

Kuehne, R. A. 1962. A classification of streams, illustrated by fish distribution in an eastern Kentucky creek. *Ecological Society of America* 43:608-614.

Kuehne, R. A. and R. W. Barbour. 1983. *The American darters*. The University of Kentucky Press, Lexington, Kentucky. 177 pp.

Kuehne, R. A. and R. M. Bailey 1961. Stream capture and the distribution of the percid fish *Etheostoma sagitta*, with geologic and taxonomic considerations. *Copeia* 1961:1-8.

Lotrich, V. A. 1973. Growth, production, and community composition of fishes inhabiting a first-, second-, and third-order stream of eastern Kentucky. *Ecological Monographs* 43:377-397.

- Lowe, J. E. 1979. The life history, behavior, and ecology of *Etheostoma sagitta* (Jordan and Swain). Unpublished M.S. thesis, The University of Tennessee. 73 pp.
- MacKenzie, D.I., J.D. Nichols, G.B. Lachman, S. Droege, J.A. Royle, and C.A. Langtimm. 2002. Estimating site occupancy rates when detection probabilities are less than one. *Ecology* 83:2248-2255.
- MacKenzie, D.I., J.A. Royle, J.A. Brown, and J.D. Nichols. 2004. Occupancy estimation and modeling for rare and elusive populations. Pages 149-172. In W.L. Thompson (Ed.). *Sampling Rare or Elusive Species: Concepts, Designs, and Techniques for Estimating Population Parameters*. Island Press, Washington, DC. 429 pp.
- Mattingly, H.T., J. E. Detar, B. K. Jones, and C. F. Walton. 2005. Factors affecting the distribution and recovery of the threatened blackside dace in Kentucky and Tennessee. Unpublished final project report prepared for the U.S. Fish and Wildlife Service, Tennessee Ecological Services Field Office. Cookeville, Tennessee. 126 pp.
- Maurer, E. P., L. Brekke, T. Pruitt, and P. B. Duffy (2007). Fine-resolution climate projections enhance regional climate change impact studies. *EOS Transactions American Geophysical Union* 88(47):504.
- McLaughlin, J.F., J.J. Hellmann, C.L. Boggs, and P.R. Ehrlich. 2002. Climate change hastens population extinctions. *Proceedings of the National Academy of Sciences of the United States of America* 99(9):6070-6074.
- Meyer, J. L. and A. B. Sutherland. 2005. Effects of excessive sediment on stress, growth, and reproduction of two southern Appalachian minnows, *Erimonax monachus* and *Cyprinella galactura*. Unpublished final project report for grant #1434-HQ-97-RU-01551, Department of Interior. University of Georgia, Institute of Ecology. Athens, Georgia.
- Osterkamp, W.R., Carey, W.P., Hupp, C.R., and Bryan, B.A. 1984. Movement of tractive sediment from disturbed lands: Specialty Conference, Coeur d'Alene, Idaho, 1984, American Society of Civil Engineers Proceedings, Hydraulic Division, p. 59-63.
- Page, L. M. 1983. *Handbook of darters*. T. F. H. Publications, Inc., Neptune City, New Jersey. 271 pp.
- Page, L. M. and B. M. Burr. 2011. *Peterson field guide to freshwater fishes of North America north of Mexico* (second edition). Houghton Mifflin Harcourt, Chicago, Illinois. 663 pp.
- Page, L. M. and G. S. Whitt. 1973. Lactate dehydrogenase isozymes, malate dehydrogenase isozymes and tetrazolium oxidase mobilities of darters (*Etheostomatini*). *Comparative Biochemistry and Physiology* 44B:611-623.
- Palmer, M. A., E. S. Bernhardt, W. H. Schlesinger, K. N. Eshleman, E. Foufoula-Georgiou, M. S. Hendryx, A. D. Lemly, G. E. Likens, O. L. Loucks, M. E. Power, P. S. White, and P. R. Wilcock. 2010. Mountaintop mining consequences. *Science* 327:148-149.
- Parker, R.S., and W. P. Carey. 1980. The quality of water discharging from the New River and Clear Fork basins, Tennessee: U.S. Geological Survey Water-Resources Investigations Report 80-37.
- Pond, G. J. 2004. Effects of surface mining and residential land use on headwater stream biotic integrity in the eastern Kentucky coalfield region. Kentucky Department for Environmental Protection, Division of Water, Frankfort, Kentucky. 48 pp.
- Pond, G. J., M. E. Passmore, F. A. Borsuk, L. Reynolds, and C. J. Rose. 2008. Downstream effects of

mountaintop coal mining: comparing biological conditions using family- and genus-level macroinvertebrate bioassessment tools. *Journal of the North American Benthological Society*. 27:717-737.

Ray, J. M. and P. A. Ceas. 2003. Fish survey and bioassessment of Station Camp Creek and Sturgeon Creek in the upper Kentucky River drainage. *Southeastern Fishes Council Proceedings* 46: 1-15.

Ruble, C., P. Rakes, M. Petty, and J. R. Shute. 2010. Captive propagation and reintroduction of the Kentucky arrow darter, *Etheostoma sagitta spilotum*, in the upper Kentucky River drainage of Kentucky. Conservation Fisheries, Inc. Interim performance report submitted to Kentucky Department of Fish and Wildlife Resources, Frankfort, Kentucky. 8 pp.

Soule, M.E. 1980. Threshold for survival: maintaining fitness and evolutionary potential. Pages 151-169 In M.E. Soule and B.A. Wilcox, eds. *Conservation biology*. Sinauer Associates, Inc., Sunderland, Massachusetts.

Thomas, C. D., A. Cameron, R.E. Green, M. Bakkenes, L.J. Beaumont, Y.C. Collingham, B.F.N. Erasmus, M Ferreira de Siqueira, A. Grainger, L. Hannah, L. Hughes, B. Huntley, A.S. van Jaarsveld, G.F. Midgley, L. Miles, M.A. Ortega-Huerta, A.T. Peterson, O.L. Phillips, and S.E. Williams. 2004. Extinction risk from climate change. *Nature* 427:145-148.

Thomas, M. R. 2008. Distribution and status of the Kentucky arrow darter (*Etheostoma sagitta spilotum*). Kentucky Department of Fish and Wildlife Resources. State and Tribal Wildlife Grant T-9-RSI-1. 9 pp. + figures.

Thomas, M. R. 2010. Personal communication regarding the taxonomic status of the Kentucky arrow darter. Ichthyologist, Kentucky Department of Fish and Wildlife Resources, Frankfort, Kentucky.

Thomas, M. R. 2013. Personal communication (email communication) regarding discovery of a new Kentucky arrow darter population in Leslie County, Kentucky. Ichthyologist, Kentucky Department of Fish and Wildlife Resources, Frankfort, Kentucky.

Thomas, M. R. and S. L. Brandt. 2012. Kentucky Arrow Darter (*Etheostoma spilotum*) reintroduction in Sugar Creek, Red Bird River Drainage, Kentucky: 2009-2012 monitoring results. Kentucky Department of Fish and Wildlife Resources. State and Tribal Wildlife Grant T-60-R. 11 pp.

Thomas, M. R. and R. B. Johansen. 2008. Systematics and conservation status of the arrow darter, *Etheostoma sagitta* (Percidae: Litocara). Page 46 In *Southeastern Fishes Council Annual Meeting Abstracts (Contributed Oral Presentation)*, Chattanooga, Tennessee. 70 pp.

U. S. Environmental Protection Agency (USEPA). 2003. Affected environment and consequences of mountaintop/valley fill coal mining. Chapter III of draft programmatic environmental impact statement. Region 3, US EPA, Philadelphia, Pennsylvania. 318 pp. (Available from: <http://www.epa.gov/region3/mtntop/index.htm>).

U. S. Fish and Wildlife Service (USFWS). 2009. Project report: Kentucky arrow darter surveys in eastern Kentucky. Kentucky Ecological Services Field Office. Frankfort, Kentucky. 4 pp.

U. S. Fish and Wildlife Service (USFWS). 2010. Project report: Kentucky arrow darter surveys in eastern Kentucky. Kentucky Ecological Services Field Office. Frankfort, Kentucky. 13 pp.

U. S. Geological Survey (USGS). 2001. Reconnaissance of stream geomorphology, low streamflow, and stream temperature in the mountaintop coal-mining region, southern West Virginia, 1999-2000. *Water-Resources Investigations Report* 01-4021.

Waters, T. F. 1995. Sediment in streams: sources, biological effects, and control. American Fisheries Society Monograph 7, Bethesda, Maryland.

Wood, P. J and P. D. Armitage. 1997. Biological effects of fine sediment in the lotic environment. Environmental Management 21:203-217.

Approval/Concurrence:

Lead Regions must obtain written concurrence from all other Regions within the range of the species before recommending changes, including elevations or removals from candidate status and listing priority changes; the Regional Director must approve all such recommendations. The Director must concur on all resubmitted 12-month petition findings, additions or removal of species from candidate status, and listing priority changes.

Approve:



07/15/2013

Date

Concur:



10/28/2013

Date

Did not concur:

_____ Date

Director's Remarks: